

## STIM1 regulates calcium signaling in taste bud cells and preference for fat in mice

Gado Dramane, ... , Philippe Besnard, Naim Akhtar Khan

*J Clin Invest.* 2012;122(6):2267-2282. <https://doi.org/10.1172/JCI59953>.

Research Article

Metabolism

Understanding the mechanisms underlying oro-gustatory detection of dietary fat is critical for the prevention and treatment of obesity. The lipid-binding glycoprotein CD36, which is expressed by circumvallate papillae (CVP) of the mouse tongue, has been implicated in oro-gustatory perception of dietary lipids. Here, we demonstrate that stromal interaction molecule 1 (STIM1), a sensor of  $\text{Ca}^{2+}$  depletion in the endoplasmic reticulum, mediates fatty acid-induced  $\text{Ca}^{2+}$  signaling in the mouse tongue and fat preference. We showed that linoleic acid (LA) induced the production of arachidonic acid (AA) and lysophosphatidylcholine (Lyso-PC) by activating multiple phospholipase  $\text{A}_2$  isoforms via CD36. This activation triggered  $\text{Ca}^{2+}$  influx in CD36-positive taste bud cells (TBCs) purified from mouse CVP. LA also induced the production of  $\text{Ca}^{2+}$  influx factor (CIF). STIM1 was found to regulate LA-induced CIF production and the opening of multiple store-operated  $\text{Ca}^{2+}$  (SOC) channels. Furthermore, CD36-positive TBCs from *Stim1*<sup>-/-</sup> mice failed to release serotonin, and *Stim1*<sup>-/-</sup> mice lost the spontaneous preference for fat that was observed in wild-type animals. Our results suggest that fatty acid-induced  $\text{Ca}^{2+}$  signaling, regulated by STIM1 via CD36, might be implicated in oro-gustatory perception of dietary lipids and the spontaneous preference for fat.

Find the latest version:

<https://jci.me/59953/pdf>



# STIM1 regulates calcium signaling in taste bud cells and preference for fat in mice

Gado Dramane,<sup>1,2</sup> Souleymane Abdoul-Azize,<sup>1,3</sup> Aziz Hichami,<sup>1</sup> Timo Vögtle,<sup>4</sup> Simon Akpona,<sup>2</sup> Christophe Chouabe,<sup>5</sup> Hassimi Sadou,<sup>3</sup> Bernhard Nieswandt,<sup>4</sup> Philippe Besnard,<sup>6</sup> and Naim Akhtar Khan<sup>1</sup>

<sup>1</sup>Unité Propre de L'Enseignement Supérieure Lipides and Signalisation Cellulaire Equipe d'Accueil 4183, Faculté des sciences de la vie, terre et environnement, Dijon, France. <sup>2</sup>Laboratoire de Biochimie, Université de Parakou, Parakou, Benin. <sup>3</sup>Laboratoire de Nutrition, Université Abdou Moumouni, Niamey, Niger. <sup>4</sup>Vascular Medicine, University Hospital, and Rudolf-Virchow-Center, DFG Research Center for Experimental Biomedicine, University of Würzburg, Würzburg, Germany. <sup>5</sup>Centre de Génétique et de Physiologie Moléculaire et Cellulaire, UMR CNRS 5534, Université Claude Bernard Lyon 1, Villeurbanne, France. <sup>6</sup>Physiologie de la Nutrition and Toxicologie, UMR U866 INSERM/Université de Bourgogne/Agro-Sup, Dijon, France.

**Understanding the mechanisms underlying oro-gustatory detection of dietary fat is critical for the prevention and treatment of obesity. The lipid-binding glycoprotein CD36, which is expressed by circumvallate papillae (CVP) of the mouse tongue, has been implicated in oro-gustatory perception of dietary lipids. Here, we demonstrate that stromal interaction molecule 1 (STIM1), a sensor of Ca<sup>2+</sup> depletion in the endoplasmic reticulum, mediates fatty acid-induced Ca<sup>2+</sup> signaling in the mouse tongue and fat preference. We showed that linoleic acid (LA) induced the production of arachidonic acid (AA) and lysophosphatidylcholine (Lyso-PC) by activating multiple phospholipase A<sub>2</sub> isoforms via CD36. This activation triggered Ca<sup>2+</sup> influx in CD36-positive taste bud cells (TBCs) purified from mouse CVP. LA also induced the production of Ca<sup>2+</sup> influx factor (CIF). STIM1 was found to regulate LA-induced CIF production and the opening of multiple store-operated Ca<sup>2+</sup> (SOC) channels. Furthermore, CD36-positive TBCs from *Stim1*<sup>-/-</sup> mice failed to release serotonin, and *Stim1*<sup>-/-</sup> mice lost the spontaneous preference for fat that was observed in wild-type animals. Our results suggest that fatty acid-induced Ca<sup>2+</sup> signaling, regulated by STIM1 via CD36, might be implicated in oro-gustatory perception of dietary lipids and the spontaneous preference for fat.**

## Introduction

The sense of taste informs the organism about the quality of ingested food. Five basic taste modalities, e.g., sweet, sour, bitter, salty, and umami, have so far been identified (1). Recent compelling evidence from rodents raises the possibility of an additional sixth taste modality devoted to the perception of lipids (2–4) that are released by the action of lingual lipases, present in the salivary secretions (5). Fukuwatari et al. (6) and Laugerette et al. (2) have documented the expression of the receptor-like lipid-binding protein CD36 in rat and mouse circumvallate papillae (CVP), respectively. Laugerette et al. (2) have provided the first evidence for the involvement of lingual CD36 in dietary lipid perception in the mouse. Indeed, *Cd36* gene inactivation completely abolished the spontaneous preference for long-chain fatty acids (LCFAs) observed in wild-type mice. This effect is lipid specific, since the preference for sweet and aversion to bitterness reported in controls were not altered in *Cd36*-null mice. These observations regarding the implication of CD36 in the perception of lipid taste have also been confirmed by other investigators (7).


The coupling of CD36 cell signaling to a downstream second messenger cascade has been explored in mouse taste bud cells (TBCs) (3, 8). The experiments conducted on CD36-positive TBCs purified from mouse CVP demonstrated that linoleic acid (LA), an LCFA, induced increases in free intracellular calcium concentrations, [Ca<sup>2+</sup>]<sub>i</sub>, by binding to CD36 (3). We have reported that LCFAs induced the production of inositol-1,4,5-trisphosphate (IP<sub>3</sub>) and, consequently, recruited Ca<sup>2+</sup> from the endoplasmic reticulum, followed by Ca<sup>2+</sup> influx via opening of Ca<sup>2+</sup> channels in these cells

(8). The binding of LCFAs to CD36-positive TBCs also induced phosphorylation of Src protein tyrosine kinases (Src-PTK), particularly of Fyn<sup>59</sup> and Yes<sup>62</sup>. In CD36-positive TBCs, LCFA-mediated increases in [Ca<sup>2+</sup>]<sub>i</sub> induced exocytosis of serotonin (5-HT) and norepinephrine into the extracellular medium. However, the mechanisms of action of the LCFAs on the opening of Ca<sup>2+</sup> channels remained unexplored (4, 9).

It is possible that activation of CD36-positive TBCs might be associated with the release of lipid messengers such as arachidonic acid (AA) that would participate in downstream signaling mechanisms. It has been shown that AA modulates activation of K<sup>+</sup> channels in isolated TBCs (10) and affects the gustatory nerve responses to taste stimuli (11). AA is known to be principally released from the *sn*-2 position of plasma membrane phospholipids by the activation of phospholipase A<sub>2</sub> (PLA<sub>2</sub>). On the basis of structural and functional properties, these enzymes have been divided into three families: secretory PLA<sub>2</sub> (sPLA<sub>2</sub>: Ib, IIa–f, III, V, X, XII), cytosolic PLA<sub>2</sub> (cPLA<sub>2</sub>: IVa–d), and Ca<sup>2+</sup>-independent PLA<sub>2</sub> (iPLA<sub>2</sub>: VIa, VIβ) (12, 13). By employing in situ hybridization techniques, Oike et al. (14) have shown that group IIa sPLA<sub>2</sub> is expressed in rat TBCs, which possess exocytotic machinery for the release of neurotransmitters. These investigators further showed that sPLA<sub>2</sub> IIa was coexpressed with transient receptor potential melastatin-5 (TRPM5) channels in these cells (14). It is also noteworthy that gustatory nerve responses are influenced by a PLA<sub>2</sub> inhibitor or by the addition of AA (11). Chorda tympani nerve responses to taste stimuli are modulated by the application of a PLA<sub>2</sub> inhibitor to the surface of the gerbil tongue (11). Therefore, some PLA<sub>2</sub> species probably function during the process of intramolecular taste signaling that is followed by the gustatory nerve response. However, there is a paucity of information on the identification and characterization of different isoforms of PLA<sub>2</sub>s in the mouse lingual

**Conflict of interest:** The authors have declared that no conflict of interest exists.

**Citation for this article:** *J Clin Invest.* 2012;122(6):2267–2282. doi:10.1172/JCI59953.



**Figure 1**

Ca<sup>2+</sup> signaling and expression of PLA<sub>2</sub> in CD36-positive TBCs. The CD36-positive TBCs and CD36-negative cells were isolated from CVP of C57BL/6J mice as described in Methods. (A) Ca<sup>2+</sup> imaging studies were performed on CD36-positive TBCs in Ca<sup>2+</sup>-free medium. The changes in intracellular Ca<sup>2+</sup> ( $F_{340}/F_{380}$ ) were monitored under the Nikon microscope (TIU) by using S Fluor 40× oil immersion objectives as described in Methods. The arrows indicate when the test molecules (LA at 20 μM; TG at 5 μM) were added into the cuvette without interruptions in the recording. (B) Total RNA isolated from CD36-positive TBCs and CD36-negative cells was analyzed with semiquantitative RT-PCR. PCR products were visualized by UV transillumination after loading into agarose gel. The lanes were run on the same gel but were noncontiguous. (C) Relative expression of PLA<sub>2</sub> mRNA in CD36-positive TBCs versus CD36-negative cells was assessed ( $n = 6$ ). \* $P < 0.001$  compared with CD36-negative cells. (D) PLA<sub>2</sub> isoforms were also detected in CD36-positive TBCs by confocal microscopy (original magnification, ×10).


TBC. Keeping in view the above-mentioned data, it was thought worthwhile to undertake the present study to characterize different isoforms of PLA<sub>2</sub>s, to demonstrate the implication of their products such as free AA and lysophospholipids in Ca<sup>2+</sup> signaling in CVP CD36-positive TBCs, and to explore the related consequence on the oro-sensory detection of dietary lipids in the mouse.

**Results**

*LA induces increases in [Ca<sup>2+</sup>]<sub>i</sub> and activates different PLA<sub>2</sub> isoforms.* Unless mentioned otherwise, all of the experiments in the present study were conducted on CD36-positive TBCs. To study the transient increase in [Ca<sup>2+</sup>]<sub>i</sub>, we used a Ca<sup>2+</sup> imaging technique. We observed

that LA induced a rapid rise in [Ca<sup>2+</sup>]<sub>i</sub> in CD36-positive TBCs (Figure 1A). Thapsigargin (TG), which blocks sarcoplasmic-endoplasmic Ca<sup>2+</sup>-ATPase (SERCA), which also evoked a rapid increase in [Ca<sup>2+</sup>]<sub>i</sub> (Figure 1A). The LA-induced increases in [Ca<sup>2+</sup>]<sub>i</sub> were greater in Ca<sup>2+</sup> buffer than that in Ca<sup>2+</sup>-free buffer (1,200 ± 50 nM in Ca<sup>2+</sup>-containing buffer vs. 300 ± 20 nM in 0% Ca<sup>2+</sup>-free buffer,  $P < 0.001$ ,  $n = 8$ ).

Since free fatty acids might be implicated in Ca<sup>2+</sup> signaling, the role of PLA<sub>2</sub> was explored. We noticed that the isolated CD36-positive and -negative cells expressed mRNA encoding three PLA<sub>2</sub> families, i.e., sPLA<sub>2</sub>, cPLA<sub>2</sub>, and iPLA<sub>2</sub> (Figure 1B). However, the CD36-positive TBCs preferentially expressed mRNA for sPLA<sub>2</sub> IIa, sPLA<sub>2</sub> V, cPLA<sub>2</sub> IVc, and iPLA<sub>2</sub> VIβ as compared with CD36-



**Figure 2**

PLA<sub>2</sub> activity in CD36-positive TBCs. (A and B) The CD36-positive TBCs ( $3 \times 10^6$ ) were treated as described in Methods. The cells were incubated with or without SSO (50  $\mu$ M) for 20 minutes and then treated with LA. For analysis of Ca<sup>2+</sup> dependence of PLA<sub>2</sub>, the Ca<sup>2+</sup>-containing medium contained CaCl<sub>2</sub> (1.2 mM), whereas Ca<sup>2+</sup>-free medium contained EGTA (2 mM) in place of CaCl<sub>2</sub>. PLA<sub>2</sub> enzyme activity was calculated as described in the kit protocol. \* $P < 0.001$  compared with unstimulated cells (basal);  $^{\S}P < 0.001$  compared with LA-treated cells. (C) The TBCs ( $2 \times 10^6$ ) were labeled for 2 hours with 1.5  $\mu$ Ci <sup>3</sup>H-AA and treated or not with PMA (10 nM) plus ionomycin (Iono, 200 nM), TG (5  $\mu$ M), or LA (20  $\mu$ M). After washing, the release of <sup>3</sup>H-AA was assessed. (D) Effect of PLA<sub>2</sub> inhibitors (15  $\mu$ M BEL, ATK, BPB, and Ar-A for 10 minutes) on LA- and TG-induced release of <sup>3</sup>H-AA in TBCs. Data (mean  $\pm$  SEM) are expressed as percentage of unstimulated cells; \* $P < 0.001$  compared with respective controls. (E) Western blot detection of 4 isoforms of PLA<sub>2</sub> in TBCs after transfection with siRNA (sPLA<sub>2</sub> IIa, cPLA<sub>2</sub> IVc, sPLA<sub>2</sub> V, and iPLA<sub>2</sub> VI $\beta$ ). NsiRNA, nontargeting siRNA. Lanes were run on the same gel but were noncontiguous, as indicated by the white line. (F) Release of <sup>3</sup>H-AA after transfection of TBCs with the siRNA isoforms mentioned in E. Experiments were performed as in D. Data (mean  $\pm$  SEM) are expressed as percentage of LA- or TG-stimulated cells; \* $P < 0.001$  compared with respective controls.

negative TBCs (Figure 1C). Figure 1D shows the expression of sPLA<sub>2</sub>, cPLA<sub>2</sub>, and iPLA<sub>2</sub> at the protein level in CD36-positive TBCs. The three isoforms seem to be distributed in the cytoplasm of resting TBCs.

LA induced the activity of the three families of PLA<sub>2</sub> in CD36-positive TBCs (Figure 2A). In order to determine whether LA evokes enzyme activity via binding to CD36, we employed sulfo-*N*-succinimidyl-oleate (SSO). SSO is known to bind specifically to CD36 (15) and, thereby, blocks the binding of the fatty acids. We observed that SSO significantly diminished the enzymatic activity of the three families of PLA<sub>2</sub> (Figure 2A). LA failed to induce the activation of the three families of PLA<sub>2</sub> in CD36-negative TBCs, and, similarly, SSO induced no actions in these cells (Figure 2A). LA failed to induce any enzyme activity of sPLA<sub>2</sub> and cPLA<sub>2</sub> in Ca<sup>2+</sup>-free medium (Figure 2B). Interestingly, LA-induced enzyme activity of iPLA<sub>2</sub> was increased several fold in Ca<sup>2+</sup>-free medium (Figure 2B).

Since AA is the main fatty acid present on the *sn*-2 position of plasma membrane phospholipids, CD36-positive TBCs were loaded with <sup>3</sup>H-AA. The incorporation of <sup>3</sup>H-AA into phospholipids of CD36-positive TBCs was as follows (dpm  $\pm$  SEM/ $2 \times 10^6$  cells): phosphatidylcholine (PC; 2,938  $\pm$  146), phosphatidylethanolamine (PE; 1,700  $\pm$  85), phosphatidylinositol/phosphatidylserine (PI/PS; 400  $\pm$  20), and neutral lipids (200  $\pm$  10). LA induced the release of <sup>3</sup>H-AA into the extracellular medium (Figure 2C). The LA-mediated effect was dependent on CD36, since SSO significantly curtailed the release of <sup>3</sup>H-AA (Table 1). Moreover, LA-induced release of <sup>3</sup>H-AA was not influenced by indomethacin (Indo), an inhibitor of the cyclooxygenase pathway; nordihydroguaiaretic acid (NDGA), an inhibitor of the lipoxygenase pathway; or 1-aminobenzotriazole (1-ABT), a cytochrome P450 epoxygenases (CYP450) inhibitor (Table 1). PMA in combination with ionomycin (Iono) and TG, agents known to induce an



**Table 1**  
Effect of different inhibitors on LA-induced release of  $^3\text{H-AA}$  in CD36-positive TBCs

Agents	Release of $^3\text{H-AA}$ (% of unstimulated cells)
LA	3,000 $\pm$ 150
LA + SSO	100 $\pm$ 20 <sup>A</sup>
LA + NDGA	2,800 $\pm$ 50 <sup>B</sup>
LA + Indo	2,950 $\pm$ 50 <sup>B</sup>
LA + 1-ABT	2,900 $\pm$ 60 <sup>B</sup>

CD36-positive TBCs loaded with  $^3\text{H-AA}$  were treated with LA containing (or not) SSO (50  $\mu\text{M}$ ), Indo (10  $\mu\text{M}$ ), NDGA (10  $\mu\text{M}$ ), or 1-ABT (10  $\mu\text{M}$ ) as mentioned in Figure 2 and Methods. Results are expressed as mean  $\pm$  SEM ( $n = 4$ ). <sup>A</sup> $P < 0.001$  compared with LA-treated cells. <sup>B</sup>NS compared with LA-treated cells.

increase in  $[\text{Ca}^{2+}]_i$  also evoked the release of  $^3\text{H-AA}$  (Figure 2C). Interestingly, LA and TG induced the release of  $^3\text{H-AA}$  to the same order of magnitude in CD36-positive TBCs (Figure 2C).

In order to shed light on the roles of different PLA<sub>2</sub>s, we used chemical inhibitors. Aristolochic acid (Ar-A) and 4-bromophenacyl bromide (BPB) inhibit sPLA<sub>2</sub> (Ib/V), and arachidonyl trifluoromethyl ketone (ATK) and bromoenol lactone (BEL) are the respective inhibitors of cPLA<sub>2</sub> IV and iPLA<sub>2</sub> VI (12). Interestingly, all of the PLA<sub>2</sub> inhibitors inhibited the TG- and LA-induced release of  $^3\text{H-AA}$  into the extracellular medium (Figure 2D).

The chemical inhibitors of PLA<sub>2</sub> are not very specific as far as their inhibitory actions are concerned. Hence, we employed siRNA technology in order to silence specifically the expression of 4 isoforms of PLA<sub>2</sub>, i.e., IIa, IVc, V, and VI $\beta$ , which are highly expressed in CD36-positive TBCs. The transfection of these cells with the siRNA diminished the expression of corresponding isoforms of PLA<sub>2</sub> (Figure 2E) and the release of  $^3\text{H-AA}$  into the extracellular medium (Figure 2F).

*PLA<sub>2</sub>-cleaved AA facilitates Ca<sup>2+</sup> influx.* Since TG selectively depletes the intracellular Ca<sup>2+</sup> store, we used this compound for further studies on Ca<sup>2+</sup> signaling. In the Ca<sup>2+</sup>-free/Ca<sup>2+</sup> reintroduction (CFCR) protocol, addition of AA and then CaCl<sub>2</sub> to Ca<sup>2+</sup>-free medium potentiated the TG-induced store-operated Ca<sup>2+</sup> (SOC) influx in a dose-dependent manner (Figure 3A). The action of AA on Ca<sup>2+</sup> entry was not influenced by Indo, 1-ABT (Figure 3A), or NDGA (data not shown). Furthermore, PLA<sub>2</sub> inhibitors were employed in order to assess their effects on TG-induced Ca<sup>2+</sup> influx. Prior addition of ATK and BEL inhibited the TG-stimulated Ca<sup>2+</sup> entry. Interestingly, Ar-A exerted the highest inhibitory effect on TG-induced Ca<sup>2+</sup> influx in CD36-positive TBCs (Figure 3B). The order of magnitude of the inhibitory effects of different PLA<sub>2</sub> inhibitors on Ca<sup>2+</sup> influx in the CFRC measurements was as follows: Ar-A > BEL > ATK (Figure 3B). The transfection of CD36-positive TBCs with siRNA of 4 isoforms of PLA<sub>2</sub> (IIa, IVc, V, and VI $\beta$ ) significantly diminished the TG-induced Ca<sup>2+</sup> influx in the CFRC protocol (Figure 3C).

*Lyso-PC and AA exert additive effects on Ca<sup>2+</sup> entry.* In Ca<sup>2+</sup>-free medium, addition of Lyso-PC did not evoke any increase in  $[\text{Ca}^{2+}]_i$ . When CaCl<sub>2</sub> was added into the cuvette, this agonist induced a significant Ca<sup>2+</sup> influx in CD36-positive TBCs (Figure 4A). Interestingly, when Lyso-PC was added after AA (Figure 4B) or when AA was added after Lyso-PC (Figure 4C), the effect on Ca<sup>2+</sup> entry was additive. PLA<sub>2</sub>-activating peptide (PLAP) was used in order to


enrich the plasma membrane of CD36-positive TBCs with lysophospholipids. We observed that PLAP evoked Ca<sup>2+</sup> signaling in these cells. 2-Aminoethoxy-diphenyl-borate (2-APB), an inhibitor of SOC channels, significantly diminished the responses evoked by both Lyso-PC and PLAP (Figure 4D).

*LA induces the production of Ca<sup>2+</sup> influx factor, which evokes Ca<sup>2+</sup> entry.* The CD36-positive TBCs were treated with LA, and then Ca<sup>2+</sup> influx factor (CIF) was extracted. Injection of LA-produced CIF evoked Ca<sup>2+</sup> influx in CD36-positive TBCs (Figure 5A). CIF was also purified from TG-treated CD36-positive TBCs (termed TG-CIF). We observed that injection of TG-CIF was also potent to induce Ca<sup>2+</sup> entry in these cells (Figure 5A). Control CIF (extracted from control unstimulated cells) exerted no effect on Ca<sup>2+</sup> influx in CD36-positive TBCs (Figure 5B). Interestingly, addition of AA after CIF had an additive effect on the increases in  $[\text{Ca}^{2+}]_i$  (Figure 5B). However, addition of CIF or Lyso-PC one after the other did not induce an additive response in terms of Ca<sup>2+</sup> influx in CD36-positive cells (Figure 5, B and C).

*CIF activates I<sub>CRAC</sub> in Jurkat T cells.* To test the hypothesis that CIF purified from LA-treated CD36-positive TBCs activates Ca<sup>2+</sup>-release activated Ca<sup>2+</sup> (CRAC) channels, we used experimental procedures known to activate CRAC currents in Jurkat T cells (16). With a low concentration of Ca<sup>2+</sup> chelating buffer (0.2 mM BAPTA) and without inositol 1,4,5-triphosphate or SERCA blocker in the pipette solution, we did not record I<sub>CRAC</sub> (that is, La<sup>3+</sup>-sensitive current was absent in this condition) in Jurkat T cells (Figure 5D, open circles). However, when CIF purified from LA-treated CD36-positive TBCs was included in the pipette solution (1:15 dilution), we observed an inhibitory effect of La<sup>3+</sup> (inset) and could measure La<sup>3+</sup>-sensitive currents (filled circles). For instance, at -80 mV, the average La<sup>3+</sup>-sensitive current densities were -0.01  $\pm$  0.11 ( $n = 7$ ) and -1.27  $\pm$  0.27 pA/pF ( $n = 9$ ) in the absence and presence of CIF in the pipette solution, respectively.

*Orai1 and Orai3 channels are implicated in Ca<sup>2+</sup> entry.* RT-PCR showed that CD36-positive TBCs expressed mRNA encoding Orai1 and Orai3 channel proteins (Figure 6A). These cells also expressed mRNA for stromal interaction molecule 1 (STIM1), implicated in sensing of depletion of the ER pool (ref. 17 and Figure 6A). The CD36-positive TBCs also expressed these channel components and STIM1 at the protein level, and the transfection of these cells with siRNA downregulated their expression, as revealed by Western blot analysis (Figure 6B). We checked whether siRNA for either Orai1 or Orai3 could curtail the expression of the other. In Western blot assays, transfection of CD36-positive TBCs by either Orai1 or Orai3 did not alter the expression of the other proteins (Figure 6B). In RT-PCR studies, we also observed no inhibition of expression of Orai1 or Orai3 mRNA in Orai3- or Orai1-transfected TBCs, respectively (Figure 6C).

LA-induced increases in  $[\text{Ca}^{2+}]_i$  were diminished in CD36-positive TBCs transfected with Orai1 and Orai3 siRNA. Moreover, the LA-evoked response was further curtailed in cells that were transfected by both Orai1 and Orai3 siRNA (Figure 6D). Similarly, the AA-induced increases were lower in TBCs transfected with Orai1 and Orai3 siRNA than in those transfected only with Orai1 or Orai3 siRNA (Figure 6D). In contrast, the increase in  $[\text{Ca}^{2+}]_i$  in response to Lyso-PC in  $[\text{Ca}^{2+}]_i$  was diminished only in Orai1-transfected cells, and no difference was observed between Orai1-transfected cells and Orai1/3-transfected cells (Figure 6D). We further conducted experiments on CD36-positive TBCs isolated from Orai1<sup>-/-</sup> mice and observed that the Lyso-PC-induced increases in



**Figure 3**

Implication of AA and PLA<sub>2</sub> in Ca<sup>2+</sup> influx in CD36-positive TBCs. (A) The CD36-positive TBCs (2 × 10<sup>6</sup>/assay) were loaded with Fura-2/AM, as described in Methods. The experiments were performed in Ca<sup>2+</sup>-free medium. The arrows indicate when the test molecules, TG (5 μM), AA (0, 10, 20 μM), and CaCl<sub>2</sub> (1.5 mM), were added. The histograms show mean ± SEM of 7 independent experiments. \*P < 0.001 compared with control (CaCl<sub>2</sub>). Indo and 1-ABT were used at 10 μM with AA. (B) Effect of different PLA<sub>2</sub> inhibitors on Ca<sup>2+</sup> influx. The experiments were performed in Ca<sup>2+</sup>-free medium. The arrows indicate when the test molecules, i.e., PLA<sub>2</sub> inhibitors (15 μM), TG (5 μM), or CaCl<sub>2</sub> (1.5 mM), were added. “No inhibitor” trace shows the recording in the absence of PLA<sub>2</sub> inhibitors. The histograms show the results, obtained from the traces, as mean ± SEM (n = 7). \*P < 0.001 compared with control, CaCl<sub>2</sub>. All of these experiments were performed in a PTI spectrofluorometer. (C) Ca<sup>2+</sup> influx after transfection of CD36-positive TBCs by siRNA of different sPLA<sub>2</sub> isoforms. Experiments were performed as in B.


[Ca<sup>2+</sup>]<sub>i</sub> were completely blocked in these cells, but the AA-induced increases in [Ca<sup>2+</sup>]<sub>i</sub> were only partially diminished (Figure 6E).

*CIF modulates iPLA<sub>2</sub> activity.* We wanted to assess the mechanism of action of CIF in the activation of Ca<sup>2+</sup> influx in CD36-positive TBCs. Smani et al. (18) have demonstrated that CIF activates iPLA<sub>2</sub> activity. We observed that CIF produced by LA did not influence sPLA<sub>2</sub> or cPLA<sub>2</sub> enzyme activity; rather, it significantly increased only iPLA<sub>2</sub> activity in these cells (Figure 7A). The calmodulin (CaM) antagonist calmidazolium (CMZ) is known to specifically bind to CaM and, therefore, dissociates iPLA<sub>2</sub>, allowing an optimal activation of this enzyme (18). CMZ and also BAPTA/AM (known to chelate intracellular Ca<sup>2+</sup>) increased iPLA<sub>2</sub> activity in CD36-positive TBCs (Figure 7B).

*STIM1 is necessary for Ca<sup>2+</sup> signaling.* STIM1, a Ca<sup>2+</sup> sensor, plays a crucial role in the regulation of Ca<sup>2+</sup> signaling during ER store depletion (17). To explore the putative involvement of STIM1 in LA-induced Ca<sup>2+</sup> signaling, we conducted experiments on CD36-positive and -negative TBCs isolated from wild-type and *Stim1*<sup>-/-</sup> mice. Both LA- and TG-induced transient increases in Ca<sup>2+</sup> signaling were significantly diminished in *Stim1*<sup>-/-</sup> CD36-positive TBCs (Figure 8A). LA, but not TG, failed to induce any increases in [Ca<sup>2+</sup>]<sub>i</sub> in CD36-negative TBCs (Figure 8A).

It is noteworthy that CIF extracted from LA-stimulated *Stim1*<sup>-/-</sup> CD36-positive TBCs (termed CIF from *Stim1*<sup>-/-</sup>) failed to induce any increase in [Ca<sup>2+</sup>]<sub>i</sub> in these cells (Figure 8B). As expected, increases in [Ca<sup>2+</sup>]<sub>i</sub> induced by AA (Figure 8C) and Lyso-PC (Figure 8D) were significantly lower in CD36-positive TBCs from *Stim1*<sup>-/-</sup> mice than those from wild-type mice. In order to assess whether the TBCs from *Stim1*<sup>-/-</sup> mice exhibit altered expression of CD36 and different isoforms of PLA<sub>2</sub>, we conducted RT-PCR studies. Figure 8E shows that the expression of mRNA encoding CD36 or 4 isoforms of PLA<sub>2</sub> (IIa, IVc, V, and VIβ) was not significantly altered in CD36-positive TBCs between wild-type and *Stim1*<sup>-/-</sup> mice (Figure 8E).

*Spontaneous gustatory preference for fat is abolished in Stim1<sup>-/-</sup> mice.* Based on these findings, it was tempting to speculate that STIM1 might play a key role in the oro-sensory perception of lipids. To explore this hypothesis, we subjected wild-type and *Stim1*<sup>-/-</sup> mice to two-bottle preference tests. As expected, wild-type mice exhibited a strong preference for a 0.1% colza oil-enriched solution as compared with a control solution containing 0.3% xanthan gum, used to emulsify the fatty acid and mimic lipid texture (Figure 9A). In contrast, this spontaneous preference for fat was significantly diminished in *Stim1*<sup>-/-</sup> mice (Figure 9A).



**Figure 4**

Effect of Lyso-PC or PLAP and AA on  $Ca^{2+}$  influx in CD36-positive TBCs. (A and B) The CD36-positive TBCs ( $2 \times 10^6$ ) were loaded with the fluorescent probe Fura-2/AM. The experiments were performed in  $Ca^{2+}$ -free medium. The arrows indicate when the test molecules, i.e.,  $CaCl_2$  (1.5 mM), Lyso-PC (5  $\mu$ M), and AA (10  $\mu$ M), were added into the cuvette. (A) Effect of Lyso-PC (5  $\mu$ M) alone. (B) Additive effects of AA (10  $\mu$ M) and Lyso-PC (5  $\mu$ M). (C) Histograms (mean  $\pm$  SEM) of 4 independent experiments. (D) Cell imaging results for  $Ca^{2+}$  influx evoked by Lyso-PC (5  $\mu$ M) and PLAP (5  $\mu$ g/ml) in the presence or absence of 2-APB (30  $\mu$ M). The values are mean  $\pm$  SEM ( $n = 7$ ); \* $P < 0.001$  compared with control.


It is important to note that the STIM1-deficient mice, despite their shortened life span and growth retardation (19, 20), were lively, actively exploring their cage, and were able to survive independent of their mothers. Only about 2 days before death did they show signs of illness. The only notable differences in STIM1-deficient mice that were monitored daily in our experiments were the smaller size and the previously described lymphoproliferative phenotype, including splenomegaly (21). To exclude that the enlarged spleen size had an impact on taste preference, we induced splenomegaly in wild-type mice by i.p. injection of  $CdCl_2$ . In these animals, the spleen size per 100 g body weight was increased by approximately 93% compared with controls. We observed no differences in spontaneous preference for fat between mice with induced splenomegaly and control mice (Figure 9B). These results suggest that it is the absence of STIM1, but not splenomegaly, that is involved in the loss of the spontaneous preference for fat. Similarly, no difference was observed in spontaneous preference for fat in mice in which splenomegaly was induced by i.v. administration of 2 doses of G-CSF at 100  $\mu$ g/kg (data not shown).

*5-HT release is abolished in TBCs from Stim1<sup>-/-</sup> mice.* We have previously shown that LCFA induced the release of neurotransmitters including 5-HT to the extracellular environment in CD36-positive

TBCs (8). We loaded the cells with  $^3H$ -5-HT and assessed its release. Figure 9C shows that LA evoked the release of  $^3H$ -5-HT into extracellular environment and SSO significantly curtailed this phenomenon. Interestingly, no exocytosis of  $^3H$ -5-HT in response to LA was observed in TBCs isolated from *Stim1<sup>-/-</sup>* mice (Figure 9C).

**Discussion**

Both CD36-positive and CD36-negative mouse TBCs expressed sPLA<sub>2</sub>, cPLA<sub>2</sub>, and iPLA<sub>2</sub>. However, CD36-positive TBCs abundantly expressed two sPLA<sub>2</sub> isoforms (IIa and V) shown to be involved in the release of AA from phospholipids in endothelial cells and macrophages (22). In fact, sPLA<sub>2</sub> V has been shown to preferentially hydrolyse PC (23). In our study, exogenous AA was also incorporated principally into PC. These gustatory cells also express cPLA<sub>2</sub> IVc, which was found to rapidly release AA in HEK293 cells (24). The CD36-positive TBCs also expressed iPLA<sub>2</sub> VI $\beta$ , well conserved in human, rat, mouse, hamster, and drosophila (25–27). We report herein that LA induces the activation of three families of PLA<sub>2</sub>s by binding to CD36 in TBCs, corroborating a recent report demonstrating that CD36 is associated with the activation of cPLA<sub>2</sub>, and to a lesser extent of iPLA<sub>2</sub>, in CHO cells (28). Besides, in vascular endothelial cells, laminar flow has been shown to induce both

**Figure 5**

Effect of CIF, AA, and Lyso-PC on  $\text{Ca}^{2+}$  influx in CD36-positive TBCs. **(A)** Effect of CIF on  $\text{Ca}^{2+}$  influx. The experiments were performed in  $\text{Ca}^{2+}$ -free medium. CIF was purified from LA-treated (CIF) or TG-treated (TG-CIF) CD36-positive TBCs. The arrows indicate when the test molecules, i.e.,  $\text{CaCl}_2$  (1.5 mM) and CIF (50  $\mu\text{l}$ ) or CIF-TG (50  $\mu\text{l}$ ), were added. **(B and C)** Histograms of the effects of C-CIF (extracted from control untreated cells), CIF, AA, and Lyso-PC. **(D)** Activation of  $\text{La}^{3+}$ -sensitive current by CIF in Jurkat T cells. Mean  $\text{La}^{3+}$ -sensitive current density-voltage relationships recorded in the absence (open circles,  $n = 7$ ) or presence (filled circles,  $n = 9$ ) of CIF included in the pipette solution (1:15 dilution). The inset shows representative ramp membrane currents recorded in response to voltage ramps, of 50-ms duration applied from  $-100$  to  $+100$  mV in a Jurkat T cell dialyzed with the CIF-containing pipette solution in the absence (a) or presence (b) of 20  $\mu\text{M}$   $\text{La}^{3+}$ . The difference curves (a and b, in gray) characterize the  $\text{La}^{3+}$ -sensitive current.

CD36 expression and activation of  $\text{iPLA}_2$  (29). Furthermore, LA-induced activity of  $\text{sPLA}_2$  and  $\text{cPLA}_2$  was increased in the presence of  $\text{Ca}^{2+}$  in the extracellular medium, as shown elsewhere (30–32). Suh et al. (33) have also shown that LA induces increases in  $[\text{Ca}^{2+}]_i$  and  $\text{cPLA}_2$  activation in chicken hepatocytes.

It was intriguing that LA and TG induced the release of  $^3\text{H}$ -AA to the same extent in CD36-positive TBCs. TG does not seem to act directly via CD36, since SSO failed to curtail its action (data not shown). In fact, Kuda et al. (28) have shown in CHO cells that an initial rise in  $[\text{Ca}^{2+}]_i$  by TG is implicated in the membrane translocation of CD36, which was responsible for  $\text{PLA}_2$  activation and the release of  $^3\text{H}$ -AA. Similarly, chemical inhibitors and siRNA of different isoforms of  $\text{PLA}_2$  significantly inhibited the LA- and TG-induced  $^3\text{H}$ -AA release in CD36-positive TBCs, as observed in Jurkat T cells, where only chemical inhibitors were used (34). Together, these observations suggest that CD36-positive cells possess the three  $\text{PLA}_2$  isoforms actively implicated in the catalysis of AA. It is noteworthy that LA did not act via cyclooxygenase, lipoxygenase, or cytochrome P450 epoxygenase pathways on the release of  $^3\text{H}$ -AA.


Since LA and TG evoked the release of  $^3\text{H}$ -AA to the same extent, and this release was similarly inhibited by various  $\text{PLA}_2$  siRNAs, we employed only TG to assess the role of free AA on  $\text{Ca}^{2+}$  influx in order to avoid interference of this fatty acid with different second messengers that might be additionally activated. Hence, in the CFCR protocol in the presence of TG, addition of AA and then exogenous  $\text{Ca}^{2+}$  exerted dose-dependent effects on the increases in  $[\text{Ca}^{2+}]_i$  in CD36-positive TBCs. Similarly, the inhibition of  $\text{PLA}_2$  activation by chemical inhibitors and siRNA curtailed the TG-induced  $\text{Ca}^{2+}$  influx in CD36-positive TBCs. These observations suggest that AA (exogenous or endogenous) evokes  $\text{Ca}^{2+}$  influx in these cells as shown elsewhere (34, 35), and its action on  $\text{Ca}^{2+}$  influx is not influenced by its metabolites (36).

We further ascertained the actions of AA and lysophospholipids on  $\text{Ca}^{2+}$  signaling in CD36-positive TBCs. Lyso-PC induced  $\text{Ca}^{2+}$  influx in these cells. Lyso-PE and Lyso-PI induced weaker  $\text{Ca}^{2+}$  influx as compared with Lyso-PC (data not shown); we therefore continued our experiments with Lyso-PC. These results are consistent with the findings of Smani et al. (18) showing that lysophospholipids, including Lyso-PC, induced the opening of  $\text{Ca}^{2+}$  channels in mouse smooth muscle cells. Boittin et al. (37) have also demonstrated that Lyso-PC triggered  $\text{Ca}^{2+}$  entry via SOC channels in dystrophic fibers. Moreover, addition of AA after Lyso-PC and vice versa induced an additive response on  $\text{Ca}^{2+}$  influx, suggesting that the channels opened by these two agents do not overlap.

CIF was first identified in Jurkat T cells as an agent that is released from ER during store depletion and, consequently, favors  $\text{Ca}^{2+}$  influx in order to refill the intracellular pools (38–40). CIF extracted from LA-treated CD36-positive TBCs induced  $\text{Ca}^{2+}$  influx in these cells and  $I_{\text{CRAC}}$  in Jurkat T lymphocytes. Similarly, CIF purified from TG-treated CD36-positive TBCs also evoked  $\text{Ca}^{2+}$  influx. These observations suggest that CIF produced by CD36-positive TBCs is a universal influx factor (38). Interestingly, Lyso-PC and CIF are probably able to activate the same type of  $\text{Ca}^{2+}$  channels in these cells. However, the  $\text{Ca}^{2+}$  channels opened by CIF and AA might belong to two different categories.

Henceforth, we wanted to shed light on the nature of the  $\text{Ca}^{2+}$  channels that are opened by AA and Lyso-PC. The SOC channels are composed of 4 channel-forming units, encoded by *Orai* genes, that are molecular constituents of the  $I_{\text{CRAC}}$  (41, 42). CD36-positive TBCs expressed both mRNA and protein of *Orai1* and *Orai3* channels. Inhibition of *Orai1* expression by siRNA significantly curtailed Lyso-PC-evoked  $\text{Ca}^{2+}$  entry. The AA-induced  $\text{Ca}^{2+}$  influx was also reduced in these cells. Interestingly, AA-evoked, but not Lyso-PC-evoked,  $\text{Ca}^{2+}$  influx was further reduced when both *Orai1* and *Orai3* proteins were knocked down compared






**Figure 6**

Expression of STIM1, Orai1, and Orai3 proteins and implication of Orai channels in Ca<sup>2+</sup> influx in CD36-positive TBCs. (A) Expression of mRNA encoding for STIM1, Orai1, and Orai3. RT-qPCR, quantitative RT-PCR. (B) Detection of STIM1, Orai1, or Orai3 proteins by Western blotting after transfection with either scrambled RNA (control) or siRNA of STIM1, Orai1, or Orai3. (C) Expression of *Orai1* and *Orai3* mRNA in CD36-positive TBCs transfected with Orai1 or Orai3 siRNA. (D) Ca<sup>2+</sup> imaging studies were performed on CD36-positive TBCs transfected with either Orai1 siRNA or Orai3 siRNA or with both. The changes in  $F_{340}/F_{380}$  were monitored under a Nikon microscope (TiU) by using S Fluor 40 $\times$  oil immersion objectives as described in Methods. The histograms show the results obtained from the traces as mean  $\pm$  SEM ( $n = 7$ ). (E) Effect of Lyso-PC (5  $\mu$ M) and AA (10  $\mu$ M) on the increases in  $[Ca^{2+}]_i$  in CD36-positive TBCs isolated from wild-type or *Orai1*<sup>-/-</sup> mice. The experiments were performed in a PTI spectrofluorometer.

with Orai1 single knockdown. Moreover, the Ca<sup>2+</sup> influx evoked by Lyso-PC was completely diminished in CD36-positive TBCs isolated from *Orai1*<sup>-/-</sup> mice, though AA-induced Ca<sup>2+</sup> influx was significantly curtailed to a greater extent in these cells. Our observations on the opening of Orai1 channels by Lyso-PC are substantiated by the conclusions of Shuttleworth (43), who has shown that homotetrameric assembly of Orai1 form CRAC channels. Similarly, Orai1 and Orai3 proteins have been shown to form a functional AA-opened Ca<sup>2+</sup> channel pore by a heteropentameric assembly of three Orai1 subunits and two Orai3 subunits in HEK293 cells (44).

STIM1, a single-transmembrane protein, has been recently identified as a Ca<sup>2+</sup> sensor in the ER membrane. Upon Ca<sup>2+</sup> depletion, STIM1 has been shown to lose Ca<sup>2+</sup> from its EF hand and to be localized into punctate structures of the ER membrane that are in close proximity (10–25 nm) to the plasma membrane, followed by SOC entry (SOCE) activation (45, 46). In our study, CIF production was significantly reduced in LA-stimulated CD36-positive TBCs isolated from *Stim1*<sup>-/-</sup> mice. These observations are in agreement with the findings of Csutora et al. (47), who have shown, in rabbit aorta smooth muscle cells, that CIF production is tightly coupled with STIM1, since its overexpression resulted in a high

**Figure 7**

Effect of CIF, CMZ, or BAPTA on PLA<sub>2</sub> activity in CD36-positive TBCs. (A) The CD36-positive TBCs ( $3 \times 10^6$ ) were treated as described in the furnished cell lysate protocol. CIF extracts were purified as described in Methods and were used at 50  $\mu$ l. (B) Cells were preincubated for 15 minutes in the presence or absence of the following agents: CMZ (3  $\mu$ M) and BAPTA/AM (10  $\mu$ M). The histograms show the results as mean  $\pm$  SEM ( $n = 7$ ); \* $P < 0.001$  compared with basal values.

level of production of CIF. Besides, NG115 cells that do not express functional STIM1 have an intrinsic deficiency of CIF production and SOC influx (47). In our study, it is possible that an LA-induced initial rise in  $[Ca^{2+}]_i$  via IP<sub>3</sub> production might be responsible for the activation of STIM1 and, consequently, for CIF production. Combined overexpression of STIM1 and Orai1 was found to produce significant amplification of SOC influx, suggesting that SOCE activation may result from conformational coupling and a direct signal transduction from STIM1 to Orai1 (48). LA, Lyso-PC, and AA failed to induce  $Ca^{2+}$  influx in CD36-positive TBCs isolated from *Stim1*<sup>-/-</sup> mice. It has been shown that AA-regulated  $Ca^{2+}$  influx is regulated by STIM1 in HEK293 cells (43).


Since STIM1 is key in orchestrating the regulation of  $Ca^{2+}$  homeostasis (CIF production and opening of  $Ca^{2+}$  channels), we conducted a study on the spontaneous preference of lipid solution in *Stim1*<sup>-/-</sup> mice. We found that the attraction for oily solution observed in wild-type mice was completely abolished in *Stim1*<sup>-/-</sup> mice. Importantly, this behavioral change was not related to the main phenotype found in *Stim1*<sup>-/-</sup> mice, i.e., splenomegaly, since no difference in the intake of oily solution was found between control animals and mice displaying drug-induced splenomegaly. We cannot completely rule out that STIM1 deficiency in other cells besides TBCs may have influenced the preference for fat. However, it is tempting to speculate that the lower attraction for fat observed in *Stim1*<sup>-/-</sup> mice might be due to the lack of LA-mediated exocytosis of 5-HT found in CD36-positive TBCs. We have previously shown that LA-induced release of neurotransmitters including 5-HT, in a  $Ca^{2+}$ -dependent manner in mouse TBCs, might be involved in the transmission of fatty taste information from tongue to brain (8).

One of the important observations in our study is that the activation of iPLA<sub>2</sub> VI $\beta$  seems very important in the absence of calcium. Indeed, we have observed that LA-evoked activity of iPLA<sub>2</sub> was increased when  $Ca^{2+}$  was chelated by EGTA in the extracellular medium (18). We further show that chelating of intracellular calcium by BAPTA/AM resulted in several-fold increases in the activity of this enzyme in CD36-positive TBCs. Another important aspect of iPLA<sub>2</sub> is that it exists in a complexed form with CaM, which keeps it in a catalytically inactive state, and removal of CaM results in its activation. In the absence of CaM, the active site of iPLA<sub>2</sub> interacts with the CaM-binding domain, resulting in

a catalytically competent enzyme, whereas reversible disruption of this interaction through the binding of CaM abrogates this interaction, resulting in a loss of iPLA<sub>2</sub> activity. In our study, we noticed that a CaM inhibitor (CMZ) activated iPLA<sub>2</sub> activity in these cells. CIF has been shown to displace inhibitory CaM from iPLA<sub>2</sub>, resulting in its activation (18).

Consistent with the data reported herein, we propose the following working hypothesis (Figure 10). Activation of CD36-positive TBCs by LA would induce the depletion of the intracellular  $Ca^{2+}$  pool via IP<sub>3</sub> production (8) and, consequently, the production of CIF, which will in turn activate iPLA<sub>2</sub> VI $\beta$  (18). The iPLA<sub>2</sub> would then produce free AA and Lyso-PC, and the latter would open Orai1 SOC channels (18, 42).  $Ca^{2+}$  influx via SOC (Orai1) channels would bring about a large increase in  $[Ca^{2+}]_i$ , which may participate in the activation of sPLA<sub>2</sub> and cPLA<sub>2</sub>, as proposed by Bolotina (42). Besides, Chang and Parekh (49) have shown that, in RBL-1 cells, only  $Ca^{2+}$  entry through SOC channels, following depletion of intracellular stores, induces cPLA<sub>2</sub> activation, and  $Ca^{2+}$  release from the intracellular stores was ineffective in activating this enzyme even though there was a robust intracellular  $Ca^{2+}$  signal. These observations suggest that only a large increase in  $[Ca^{2+}]_i$ , contributed by  $Ca^{2+}$  influx, might participate in the activation of  $Ca^{2+}$ -dependent PLA<sub>2</sub>s (49). The significant decrease in the LA-induced activation of both sPLA<sub>2</sub> and cPLA<sub>2</sub> in  $Ca^{2+}$ -free medium in our study, strongly indicates that  $Ca^{2+}$  influx is implicated in the activation of these two families of enzymes. The massive production of Lyso-PC via the activation of sPLA<sub>2</sub> and cPLA<sub>2</sub> would again open Orai1 channels, while the generation of AA will activate the AA-activated channels formed by Orai1 and Orai3 proteins (44). This increase in  $[Ca^{2+}]_i$  might be involved in exocytosis of neurotransmitters such as 5-HT (8).

Finally, our study shows that the  $Ca^{2+}$  signaling mechanism triggered by LA in CD36-positive TBCs is under the control of STIM1, which is also necessary for oro-perception of dietary lipids. However, further studies are required to explore in-depth the intermediate molecules between STIM1 and CIF production. It is not well understood how Lyso-PC (or its derived products) opens  $Ca^{2+}$  channels in a membrane-delimited manner. The involvement of intermediate steps and additional components responsible for signal transduction from STIM1 in the ER membrane to channel proteins in plasma membrane remains to be explored.



**Figure 8**


Effect of STIM1 deficiency on Ca<sup>2+</sup> signaling. Ca<sup>2+</sup> imaging studies were performed on CD36-positive TBCs isolated from wild-type or *Stim1*<sup>-/-</sup> mice. The changes in  $F_{340}/F_{380}$  were monitored under a Nikon microscope (TiU) by using S Fluor 40× oil immersion objectives as described in Methods. **(A)** Effects of LA (20 μM) and TG (5 μM) in CD36-positive or CD36-negative TBCs. **(B)** Effect of CIF extracted from LA-stimulated TBCs isolated from wild-type or *Stim1*<sup>-/-</sup> mice (CIF/ *Stim1*<sup>-/-</sup>) on Ca<sup>2+</sup> signaling. **(C and D)** Effects of AA (10 μM) and Lyso-PC (5 μM) on increases in [Ca<sup>2+</sup>]<sub>i</sub>. The data in **A–D** represent mean ± SEM (*n* = 7). \**P* < 0.001 compared with TBCs isolated from wild-type mice. **(E)** Expression of mRNA encoding CD36 and different isoforms of PLA<sub>2</sub> in WT and *Stim1*<sup>-/-</sup> CD36-positive TBCs. Data represent mean ± SEM (*n* = 6). In **A** and **C**, the colored time-lapse changes in the increases in [Ca<sup>2+</sup>]<sub>i</sub> were recorded in wild-type CD36-positive cells.

The results presented herein may help understand the basic modalities involved in the oro-gustatory perception of dietary fat and reinforce the emerging concept of the existence of a sixth taste modality devoted to the perception of dietary lipids. Consistent with this hypothesis, it has been recently reported in humans that oral sensitivity to a LCFA (oleic acid) is related to fat intake and body mass index (50) and that CD36 influences oral sensitivity to fat in obese subjects (51). Therefore, a better knowledge of the

CD36 signaling cascade activated in gustatory papillae by dietary fat might lead to new nutritional or pharmacological strategies for limiting fat attraction and obesity risk.

**Methods**

**Materials.** C57BL/6J mice were obtained from Janvier Elevage. SSO was a gift from J.F. Glatz (Cardiovascular Research Institute, Maastricht, The Netherlands). The culture medium RPMI 1640 and L-glutamine were pur-

**Figure 9**

Effect of *Stim1* gene invalidation and splenomegaly on gustatory preference for lipids. (A) Wild-type and *Stim1*<sup>-/-</sup> mice were subjected to a 12-hour two-bottle preference test. Xanthan gum (0.3%) was used to emulsify colza oil (0.1%) in water and to mimic the lipid texture. \**P* < 0.001 compared with control solution in WT animals; <sup>§</sup>*P* < 0.001 between oil-fed WT and oil-fed *Stim1*<sup>-/-</sup> mice. (B) Splenomegaly was induced by i.p. administration of CdCl<sub>2</sub> as described in Methods, and mice were subjected to a 12-hour two-bottle preference test. \**P* < 0.001 compared with control solution in vehicle-administrated animals. Data represent mean ± SEM (*n* = 6). (C) Release of <sup>3</sup>H-5-HT into the extracellular environment in the presence of LA with or without SSO from CD36-positive TBCs isolated from wild-type or *Stim1*<sup>-/-</sup> mice. \**P* < 0.001 compared with control in CD36-positive TBCs isolated from WT animals; <sup>§</sup>*P* < 0.001 between LA- and LA plus SSO-treated cells in CD36-positive TBCs isolated from WT animals. Data represent mean ± SEM (*n* = 7).


chased from Lonza. Fura-2/AM was procured from Invitrogen. Elastase and dispase were purchased from Serlabo Technologies and Roche Diagnostics, respectively. Anti-PLA<sub>2</sub> antibodies and anti-CD36 antibody coupled to phycoerythrin and anti- $\alpha$ -gustducin antibody were procured from Santa Cruz Biotechnology Inc. Anti-Orai1 and anti-Orai3 antibodies were obtained from Interchim. ATK, BEL, and PLA<sub>2</sub> enzyme activity kit were bought from Cayman Chemical. Methyl arachidonyl fluorophosphonate was obtained from Calbiochem. <sup>3</sup>H-AA (specific activity, 217 Ci/mmol) and 5-hydroxytryptamine creatinine sulfate (serotonin) (5-[1,2-<sup>3</sup>H(N); specific activity, 25 Ci/mmol) were purchased from Amersham. Anti-PLA<sub>2</sub> antibodies were obtained from United States Biological. Sep-Pak Vac C18 cartridges were obtained from Millipore. All other chemicals, including AA (20:4 n-6), LA (18:2 n-6), collagenase type I, trypsin inhibitor, Ar-A, 4-bromo phenacyl bromide (BPB), and anti-STIM1 antibodies, were obtained from Sigma-Aldrich.

**Isolation of CD36-positive TBCs from mouse CVP.** CD36-positive cells from mouse lingual CPV were isolated according to previously published procedure (3, 8). Briefly, lingual epithelium was separated from connective tissues by enzymatic dissociation (elastase and dispase mixture, 2 mg/ml each, in Tyrode buffer: 120 mM NaCl; 5 mM KCl; 10 mM HEPES; 1 mM CaCl<sub>2</sub>; 1 mM MgCl<sub>2</sub>; 10 mM glucose; 10 mM Na pyruvate, pH 7.4). CD36-positive cells were isolated by incubating lingual epithelium in RPMI 1640 medium containing 2 mM EDTA, 1.2 mg/ml elastase, 0.6 mg/ml collagenase (type I), and 0.6 mg/ml trypsin inhibitor at 37°C for 10 minutes, followed by centrifugation (600 g, 10 minutes). The mixture of different cell populations was incubated with anti-CD36 antibody coupled to phycoerythrin for 2 hours, followed by a wash with PBS, pH 7.4 (600 g, 10 minutes), and resuspended in a solution containing microbeads coupled to anti-phycoerythrin IgG. The CD36-positive TBCs were isolated by passing through the MACS columns of the Miltenyi magnet system. Both the cell populations, after separation, were suspended in fresh RPMI 1640 medium containing 10% fetal calf serum, 200 U/ml penicillin, and 0.2 mg/ml streptomycin, seeded onto a BD BioCoat Poly-D-Lysine-coated dishes, and cultured for 24 hours. At the end of this period, the cells were used for the experiments or stained with trypan blue to assess their viability.

**Detection of mRNA by real-time PCR (RT-PCR).** Total RNA was extracted from CD36-negative and/or CD36-positive TBCs by using TRIzol (Invitrogen) and subjected to DNase treatment using the RNase-free DNase Set (QIAGEN). One microgram of total RNA was reverse transcribed with SuperScript II RNase H reverse transcriptase using oligo(dT) according to the manufacturer's instructions (Invitrogen). RT-PCR was performed on the iCycler iQ real-time detection system, and amplification was undertaken by using SYBR Green I detection. Oligonucleotide primers were based on the sequences of mouse genes in the GenBank database (Table 2). The amplification was carried out in a total volume of 25  $\mu$ l, which contained 12.5  $\mu$ l SYBR Green Supermix buffer (50 mM KCl; 20 mM Tris-HCl [pH 8.4]; 3 mM MgCl<sub>2</sub>; 0.2 mM of each dNTP, 0.63 U iTaq DNA polymerase, and SYBR Green 1.0 nM fluorescein) and 12.5  $\mu$ l (0.3  $\mu$ M) of each primer and diluted cDNA.

Results were evaluated by iCycler iQ software including standard curves, amplification efficiency (*E*), and cycle threshold (*Ct*). Relative quantification of mRNA in different groups was determined as follows:  $\Delta\Delta Ct = \Delta Ct$  of gene of interest -  $\Delta Ct$  of  $\beta$ -actin;  $\Delta Ct = Ct$  of CD36-positive or CD36-negative cells -  $Ct$  of total cells. Relative quantity (RQ) was calculated as follows:  $RQ = (1 + E)^{-\Delta\Delta Ct}$ . For semiquantitative determinations, amplification was performed under the same conditions as described above for 30 cycles, followed by a final extension period of 72°C for 10 minutes. Reaction products were electrophoresed on a 1.5% agarose gel impregnated with ethidium bromide. The RNA pattern was visualized by UV transillumination.

**Detection of different isoforms of PLA<sub>2</sub> by confocal microscopy.** Cytospin-prepared slides containing CD36-positive cells were fixed in 95% ethanol and rehydrated in 0.1 M PBS, pH 7.4. Slides were blocked in PBS containing 5% fetal calf serum and 0.2% Triton X-100 for 30 minutes at room temperature before overnight incubation at 4°C with rabbit anti-cPLA<sub>2</sub>, donkey anti-sPLA<sub>2</sub>, and iPLA<sub>2</sub> antibodies (1:100 dilution). After washing, slides were incubated for 2 hours at room temperature with secondary antibodies, FITC-conjugated anti-mouse IgG for the expression of sPLA<sub>2</sub> and iPLA<sub>2</sub>, and Alexa Fluor 633-conjugated anti-mouse IgG for cPLA<sub>2</sub> detection (1:600 dilutions). Staining specificity was assessed by treating slides in the



**Figure 10**  
 Schematic representation of LA-induced Ca<sup>2+</sup> signaling in CD36-positive TBCs. LA binds to CD36 and leads to IP<sub>3</sub> production (8) and depletes intracellular Ca<sup>2+</sup> stores. Upon Ca<sup>2+</sup> depletion, STIM1 will lose Ca<sup>2+</sup> from its EF hand and be localized into punctate structures of the ER membranes, which are in close proximity to the plasma membrane. STIM1 triggers production of CIF that displaces inhibitory CaM from a plasma membrane-bound Ca<sup>2+</sup>-independent phospholipase A<sub>2</sub> (iPLA<sub>2</sub>β), which produces Lyso-PC and free AA. The Lyso-PC will activate Orai1-dependent SOC channels that are responsible for SOCE. A massive Ca<sup>2+</sup> influx via these channels may activate sPLA<sub>2</sub> and cPLA<sub>2</sub> and give rise to massive production of Lyso-PC and AA that will open Orai1 and Orai1/3 channels, respectively, and a massive increase in [Ca<sup>2+</sup>]<sub>i</sub> might be involved in exocytosis of neurotransmitters such as 5-HT, responsible for gustatory perception of dietary lipids.

absence of primary antibodies. After 3 washings with PBS, a drop of Aqua Poly-Mount (Polysciences) mounting medium was added on the slide for the analysis under fluorescence microscope (Zeiss).

**Measurement of PLA<sub>2</sub> enzyme activity.** The activity of different isoforms of PLA<sub>2</sub> was determined as per a protocol furnished with the kit. In brief, the CD36-positive TBCs (3 × 10<sup>6</sup>/assay) were scraped off by a rubber policeman, sonicated in 1 ml cold buffer (50 mM HEPES, 1 mM EDTA, pH 7.4), and centrifuged at 10,000 g for 15 minutes at 4°C. The supernatant was removed and kept on ice. The amount of protein was determined by bicinchoninic acid (Pierce), and the assays were performed the same day. The assays for cPLA<sub>2</sub> and sPLA<sub>2</sub> were as per protocols; however, to detect the activity of iPLA<sub>2</sub>, phospholipase activity was assayed by incubating the samples with the substrate arachidonoyl Thio-PC for 1 hour at 20°C in a modified Ca<sup>2+</sup>-free buffer (300 mM NaCl; 0.5% Triton X-100; 60% glycerol; 4 mM EGTA; 10 mM HEPES at pH 7.4, and 2 mg/ml BSA) as suggested by Smani et al. (18). We also added the inhibitors of sPLA<sub>2</sub> (Ar-A and BPB) and cPLA<sub>2</sub> (ATK) at 15 μM to block any interference by these enzymes. The reaction was stopped by the addition of DTNB for 5 minutes, and the absorbance was determined at 405 nm using a standard plate reader.

The PLA<sub>2</sub> activity was expressed in nmol/min/mg protein. The background iPLA<sub>2</sub>-independent component of basal lipase activity was determined in control samples when all specific iPLA<sub>2</sub> activity was inhibited with BEL (15 μM). This background activity was subtracted from all readings.

**Measurement of the release of <sup>3</sup>H-AA.** The PLA<sub>2</sub> activation was also measured by the release of AA (34). In brief, CD36-positive TBCs (2 × 10<sup>6</sup>/assay) were labeled with <sup>3</sup>H-AA (1.5 μCi/3 × 10<sup>8</sup> cells) for 2 hours in RPMI 1640 serum-free medium supplemented with 0.2% fatty acid-free BSA. At the end of the incubation, cells were washed twice with RPMI 1640 serum-free medium containing 0.2% BSA. The cells were resuspended in 500 μl RPMI 1640 medium supplemented with 0.5% BSA at a final concentration of 12 × 10<sup>6</sup> cells/ml and treated with 15 μM PLA<sub>2</sub> inhibitors or vehicle (DMSO, 0.1% v/v) for 10 minutes with or without TG (5 μM) or LA (20 μM). Cells were centrifuged (1,250 g, 3 minutes) and lipids were extracted from both supernatant and pellets and separated by thin layer chromatography (TLC); the zone containing fatty acids and phospholipids was scraped into vials, and radioactivity was measured (28) by counting in a liquid scintillation analyzer (Packard). The released <sup>3</sup>H-AA was expressed as percentage of total <sup>3</sup>H-AA in cells and normalized to unstimulated cells.



**Table 2**  
Sequences of the primers used to study the expression of mRNA of different genes

Gene	Primer sequence
$\beta$ -Actin	Forward: 5'-AGAGGGAAATCGTGCCTGAC-3' Reverse: 5'-CAATAGTGATGACCTGGCCGT-3'
iPLA <sub>2</sub> VI	Forward: 5'-CCGTATGAAGGACGAGGTGT-3' Reverse: 5'-CGGTGGCTTCAGGTTAATGT-3'
iPLA <sub>2</sub> VI $\beta$	Forward: 5'-ATGATTATCAGCATGGACAGCA-3' Reverse: 5'-ACACAGGTTACAGGCACTTGAGG-3'
cPLA <sub>2</sub> IVc	Forward: 5'-TTCAGCAGGCTCTGGGTACT-3' Reverse: 5'-TGGTCGAGAACTGCTCCTT-3'
cPLA <sub>2</sub> IVf	Forward: 5'-ACGGAACCTTTGGCTCTGA-3' Reverse: 5'-GACACTGACACTCCCCTGT-3'
cPLA <sub>2</sub> IVd	Forward: 5'-AGATGAGGTCCCTGTTGTGG-3' Reverse: 5'-TTCCCGAAGCTTCAGTGTCT-3'
cPLA <sub>2</sub> IVa	Forward: 5'-GTCCCTTGCCTCTCTTCAGC-3' Reverse: 5'-TTTGCCATGCCAATCTCATA-3'
sPLA <sub>2</sub> IIa	Forward: 5'-GGCTGTGTGTCAGTCCGATAAA-3' Reverse: 5'-TTTTCAGCATTGGGCTTCT-3'
sPLA <sub>2</sub> X	Forward: 5'-GCTGTTATTGTGGCCTTGGT-3' Reverse: 5'-TTGGCATTGTTCTCTGCTG-3'
sPLA <sub>2</sub> V	Forward: 5'-GATGCACGACCGTTGTTATG-3' Reverse: 5'-CAGGCAGTAGACCAGCTTCC-3'
Stim1	Forward: 5'-GGTAGCCGAAACACACGAAT-3' Reverse: 5'-GAAAGGAAGGGAGGTGAAGG-3'
Orai1	Forward: 5'-TCCCTGGTCAGCCATAAGAC-3' Reverse: 5'-TCATGGAGAAGGGCATAAGG-3'
Orai3	Forward: 5'-GCCACCTCTGTAAGCTCTG-3' Reverse: 5'-TCCTGGAGGAGCAAACAAC-3'

The incorporated radioactivity into phospholipids was also determined. Briefly, phospholipids were extracted by chloroform/methanol/NaCl 2 M (1:1:0.9, v/v), and then phospholipid classes were separated by TLC using silica G60 and the following solvent: chloroform/methanol/acetic acid (35:14:2.7, v/v/v). Phospholipid classes comigrating with authentic standards were scraped off, and radioactivity was quantified by adding 2 ml of scintillation cocktail in a liquid scintillation analyzer (Packard).

**Measurement of Ca<sup>2+</sup> signaling.** CD36-positive TBCs, cultured for 24 hours, were washed with a buffer containing the following: 3.5 mM KH<sub>2</sub>PO<sub>4</sub>; 17.02 mM Na<sub>2</sub>HPO<sub>4</sub>; 136 mM NaCl, pH 7.4. The cells were then incubated with Fura-2/AM (1  $\mu$ M) for 60 minutes at 37°C in loading buffer that contained the following: 110 mM NaCl; 5.4 mM KCl; 25 mM NaHCO<sub>3</sub>; 0.8 mM MgCl<sub>2</sub>; 0.4 mM KH<sub>2</sub>PO<sub>4</sub>; 20 mM HEPES-Na; 0.33 mM Na<sub>2</sub>HPO<sub>4</sub>; 1.2 mM CaCl<sub>2</sub>, pH 7.4.

After loading, the cells (2  $\times$  10<sup>6</sup>/ml) were washed 3 times (600 g, 10 minutes) and remained suspended in the identical buffer. For experiments in Ca<sup>2+</sup>-free medium, CaCl<sub>2</sub> was replaced by EGTA (2 mM). [Ca<sup>2+</sup>]<sub>i</sub> was measured according to Grynkiewicz et al. (52). The fluorescence intensities were measured in the ratio mode in a PTI spectrofluorometer at 340 nm and 380 nm (excitation filters) and 510 nm (emission filter). The cells were continuously stirred throughout the experiment. The test molecules were added into the cuvettes in small volumes with no interruptions in recordings. The intracellular concentrations of free Ca<sup>2+</sup> ([Ca<sup>2+</sup>]<sub>i</sub>) were calculated by using the following equation: [Ca<sup>2+</sup>]<sub>i</sub> =  $K_d(RR_{min})/(R_{max}R)(Sf_2/Sb_2)$ , where  $K_d$  is the dissociation constant for Fura-2/Ca<sup>2+</sup> complex;  $R$  is the emission ratio with excitation at 340 nm divided by excitation at 380 nm;  $R_{min}$  is the ratio in the presence of no Ca<sup>2+</sup>;  $R_{max}$  is the ratio of saturating [Ca<sup>2+</sup>]<sub>i</sub>; and  $Sf_2/Sb_2$  is the ratio of 380 nm excitation fluorescence at zero

and saturating [Ca<sup>2+</sup>]<sub>i</sub>. A value of 224 nM for  $K_d$  was added into the calculations.  $R_{max}$  and  $R_{min}$  values were obtained by addition of ionomycin (5  $\mu$ M) and MnCl<sub>2</sub> (2 mM), respectively.

We designed a CFRCR protocol for some experiments, conducted in Ca<sup>2+</sup>-free medium, and exogenous CaCl<sub>2</sub> was added as indicated in the figures.

**Ca<sup>2+</sup> imaging studies.** The CD36-positive TBCs were cultured on WillCo-dish wells with a glass bottom and loaded with Fura-2/AM as described above. The changes in intracellular Ca<sup>2+</sup> ( $F_{340}/F_{380}$ ) were monitored under a Nikon microscope (TiU) by using an S Fluor 40 $\times$  oil immersion objective. The planes were taken at  $z$  intervals of 0.3  $\mu$ m, and NIS-Elements software was used to deconvolve the images. The microscope was equipped with an EM-CCD (Lucas) camera for real-time recording of 16-bit digital images. The dual excitation fluorescence imaging system was used for studies of individual cells. The changes in intracellular Ca<sup>2+</sup> were expressed as  $\Delta$ ratio, which was calculated as the difference relative to the peak  $F_{340}/F_{380}$  ratio. The data were summarized from the large number of individual cells (20–40 cells in a single run, with 3–9 identical experiments done in at least 3 cell preparations).

**Electrophysiological studies.** Electrophysiological recordings were carried out at room temperature (20–23°C) in the conventional whole-cell configuration of the patch-clamp technique (53). Membrane capacitance ( $C_m$ ) was determined as previously described (54). The average values for  $C_m$  in Jurkat T cells was 18.4  $\pm$  1.0 pF ( $n$  = 28). The series resistance ranged from 1.5 to 8.2 M $\Omega$ . Membrane capacitance and series resistance were not compensated. Command voltage and data acquisition were performed with pClamp software (Axon Instruments). Membrane currents were evoked every 3 seconds from a holding potential of 0 mV, by voltage ramps of 50-ms duration, applied from –100 to +100 mV, sampled at 20 kHz, and filtered at 3 kHz. Current traces were not corrected for the leak. The external solution contained (in mM): 145 NaCl; 2.8 KCl; 2 MgCl<sub>2</sub>; 20 CaCl<sub>2</sub>; 6 CsCl; 10 glucose; 10 HEPES; adjusted to pH 7.4 with NaOH. The internal (i.e., the pipette) solution contained (in mM): 145 Cs glutamate; 8 NaCl; 1 MgCl<sub>2</sub>; 4 MgATP; 0.2 BAPTA; 10 HEPES adjusted to pH 7.2 with CsOH. CIF purified from LA-treated CD36-positive TBCs was included in the pipette solution to a dilution of 1:15. The  $I_{CRAC}$  was recorded as the La<sup>3+</sup>-sensitive current obtained by subtraction of the La<sup>3+</sup>-resistant current from the total current. For the clarity of the figure showing average current ramps, only one point every 5 mV is shown.

**Table 3**  
Effect of different siRNA transfections on the viability of CD36-positive TBCs

Agents/treatments	RFU/assay
Control (scrambled RNA transfected) cells	10 $\times$ 10 <sup>6</sup> $\pm$ 0.10 $\times$ 10 <sup>6</sup>
Saponin-treated cells	50 $\times$ 10 <sup>3</sup> $\pm$ 20 $\times$ 10 <sup>3A</sup>
NsiRNA	9.9 $\pm$ 0.10 $\times$ 10 <sup>6B</sup>
sPLA <sub>2</sub> siRNA IIa	9.5 $\times$ 10 <sup>6</sup> $\pm$ 0.30 $\times$ 10 <sup>6B</sup>
sPLA <sub>2</sub> siRNA V	9.4 $\times$ 10 <sup>6</sup> $\pm$ 0.35 $\times$ 10 <sup>6B</sup>
cPLA <sub>2</sub> siRNA IVc	9.6 $\times$ 10 <sup>6</sup> $\pm$ 0.40 $\times$ 10 <sup>6B</sup>
iPLA <sub>2</sub> siRNA VI $\beta$	9.6 $\times$ 10 <sup>6</sup> $\pm$ 0.32 $\times$ 10 <sup>6B</sup>
Stim1 siRNA	9.6 $\times$ 10 <sup>6</sup> $\pm$ 0.30 $\times$ 10 <sup>6B</sup>
Orai1 siRNA	9.7 $\times$ 10 <sup>6</sup> $\pm$ 0.10 $\times$ 10 <sup>6B</sup>
Orai3 siRNA	9.8 $\times$ 10 <sup>6</sup> $\pm$ 0.10 $\times$ 10 <sup>6B</sup>

The fluorescence intensity was measured and the relative arbitrary fluorescence units (RFU) were determined in viable and saponin-treated (0.001%) cells (as negative control) (10<sup>5</sup> cells/ml). Control untreated cells showed RFU of 14  $\times$  10<sup>6</sup>  $\pm$  0.10  $\times$  10<sup>6</sup>. <sup>A</sup> $P$  < 0.001 compared with control (scrambled RNA-transfected) cells. <sup>B</sup>NS compared with control (scrambled RNA transfected) cells.



**Purification of CIF extract.** The CIF extract was purified as described by Smani et al. (18). The CD36-positive TBCs ( $10 \times 10^6$  cells/assay), treated or not with LA (20  $\mu$ M) or TG (5  $\mu$ M) for 15 minutes, were washed 3 times and resuspended in 0.85 ml HBSS. The suspension was extracted with 0.2 ml of 1 M HCl for 30 minutes at room temperature. After centrifugation, the supernatant was neutralized with 10 M NaOH, and 10 mM BaCl<sub>2</sub> was added to precipitate compounds containing vicinal phosphates (including IP<sub>3</sub>). After centrifugation, the supernatant was lyophilized, and the residue was extracted with 0.8 ml methanol with continuous mixing for 15 minutes. The methanol extract was loaded onto a Sep-Pak Vac C18 cartridge, and the cartridge was washed with 0.8 ml methanol. The combined methanol eluates were dried at 30°C under a stream of nitrogen and resuspended in 100  $\mu$ l of 0.1 M acetic acid. The reconstituted extract was clarified by centrifugal ultrafiltration through a Centricon 30 (Millipore). We used an additional step of purification of CIF by HPLC Partisil 10 SAX anion exchange 10  $\mu$ m, 250  $\times$  4.6 mm column (Keystone Scientific), and the elution was performed by a linear salt and pH gradient from 5 to 750 mM (NH<sub>4</sub>)<sub>2</sub>PO<sub>4</sub> and from pH 2.8 to 3.7, respectively (55). At concentrations as low as 1  $\mu$ l, the CIF extract induced increases in [Ca<sup>2+</sup>]<sub>i</sub> in CD36-positive TBCs and other cells.

**siRNA knockdown of STIM1, Orai1, Orai3, and PLA<sub>2</sub> isoforms.** The CD36-positive TBCs were transfected with the siRNA ON-Target plus Smart pool (25 nM) designed against Orai1, Orai3, STIM1, different PLA<sub>2</sub> isoforms, or non-targeting siRNA as a control (Dharmacon). The sequences of the STIM1/siRNA duplexes were as follows: 5'-GAAGUAGGCAGACUAGGGU-3', 5'-AAACAUAGCACCUUCCAUG-3', 5'-GAUCGGAGCCACAGGCA-GA-3', and 5'-UACAGUGGCUCAUUACGUA-3'. The sequences of the Orai1/siRNA duplexes were 5'-AAUAUUAUACAAGGGACA-3', 5'-UCA-CUAAACACCAGUCUUU-3', 5'-GGAAAUGGCUCGGGGACAA-3', and 5'-ACAUCGAGGCUGUGAGCAA-3'. The sequences of the Orai3/siRNA duplexes were 5'-GAUCCUUGGUUGCCACAA-3', 5'-GGAGAAGAUU-CAGCCUUU-3', 5'-CUGUGGGACUAGGUGUUU-3', and 5'-AGGUA-CAGCUGGAGAACGA-3'. The sequences of the PLA<sub>2</sub> siRNA/duplex were as follows: PLA<sub>2</sub> IIa, 5'-GCUUAGCUAUGCCUUCUAU-3', 5'-ACUCAUGACU-GUUGCUACA-3', 5'-GAUCAUGGCCUUUGGCUCA-3', 5'-GAACAU-UGCGCAGUUUGGG-3'; PLA<sub>2</sub> V, 5'-ACGGAACACAUGCGAUCUA-3', 5'-GAGAUUAUAUCAUGCGCGA-3', 5'-CAAGUCAUCUAUCGUUUUA-3', 5'-UUGAGAAGGUGACCGGGAA-3'; PLA<sub>2</sub> VI, 5'-GGUGCGAGA-UGGUCGGCAU-3', 5'-GCACCUGGCCAUGUCGAAA-3', 5'-GCUC-CAUGAGGGACGAGAA-3', 5'-GGGCCAAGAACGCCGAGAU-3'; PLA<sub>2</sub> IVc, 5'-CCUCUAUAAACACGGUAAA-3', 5'-GGAAUGAGACCUCGCGAGA-3', 5'-UAGAAGGACCAGUGACAUA-3', 5'-AGAGAUUUGCUGAGGGUUAU-3'. Twenty-four hours before treatment, the cells were placed in the culture medium without antibiotics and further transfected with siRNA by using DharmaFECT (Dharmacon). After 24-hour incubation, the medium was aspirated and replaced with fresh medium without siRNA for 48 hours before Ca<sup>2+</sup> measurements or Western blot analysis.

The cell viability was routinely checked by Blue Cell Viability Kit (Abnova). The assay kit utilizes the redox dye resazurin, which is transformed into a fluorescent molecule upon reduction by viable cells. The fluorescence intensity (excitation, 530 nm; emission, 600 nm) after an incubation step was measured in a plate reader, and the relative arbitrary fluorescence units (RFU) were determined, after 24 hours, in control and transfected or saponin-treated (0.001%; as negative control) cells. We observed that the transfection of CD36-positive TBCs by any of the siRNAs did not significantly reduce cell viability as compared with scrambled siRNA (Table 3). We obtained around 85% of viable cells after transfection by the Dharmacon kit (for example, control non-transfected cells,  $12 \times 10^6 \pm 0.50 \times 10^6$  RFU vs. control scrambled RNA-transfected cells,  $10 \times 10^6 \pm 0.50 \times 10^6$ ).

**Western blot detection of STIM1, Orai1, Orai3, and PLA<sub>2</sub> isoforms.** After transfection and culture for 24 hours, the CD36-positive TBCs were lysed with 50  $\mu$ l of buffer containing the following: 20 mM HEPES pH 7.3; 1 mM EDTA; 1 mM EGTA; 0.15 mM NaCl; 1% Triton X-100; 10% glycerol; 1 mM PMSF; 2 mM sodium orthovanadate and anti-protease cocktail (2  $\mu$ l in 1 ml of buffer). After centrifugation (13 000 g, 1 minute), cells lysates were used immediately or stored at -20°C. The protein content was determined by bicinchoninic acid (Pierce). Denatured proteins (30  $\mu$ g) were separated by SDS-PAGE (10%) and transferred to PVDF membranes, and immunodetection was performed by using anti-STIM1 (4  $\mu$ g/ml), anti-Orai1 (2  $\mu$ g/ml), or anti-Orai3 antibodies (4  $\mu$ g/ml). For PLA<sub>2</sub> detection, antibodies were used at a 1:2,000 dilution, and the proteins were revealed by secondary anti-goat antibodies. After treating the membranes with peroxidase-conjugated goat anti-mouse secondary antibodies, peroxidase activity was detected with ECL reagents (Amersham).

**Mice deficient in STIM1 and Orai1.** Mice homozygously deficient in *Stim1* or *Orai1* were generated from embryonic stem cell clones and germline transmission, as described elsewhere (19, 56). Because *Stim1*<sup>-/-</sup> and *Orai1*<sup>-/-</sup> mice showed growth retardation and early lethality, 3- to 5-week-old mice were used in the experiments and wild-type littermates were used as controls. There were no significant differences in the yield of CD36-positive TBCs between wild-type and *Stim1*<sup>-/-</sup> or *Orai1*<sup>-/-</sup> mice.

**Induction of splenomegaly.** Mice were injected i.p. with a single dose of cadmium chloride (CdCl<sub>2</sub>) (1.8 mg/kg body weight) as per the protocol of Pathak and Khandelwal (57). Control mice were injected with the vehicle (NaCl 9 g/l). Animals were weighed before and after the study. Seventy-two hours after administration, animals were sacrificed, and their spleens were immediately dissected out, weighed, and examined under the microscope. We observed that cadmium increased spleen weight by 93% compared with the vehicle-injected mice.

**Two-bottle preference test of lipid solution.** Spontaneous preference for lipid-enriched solutions was investigated by means of the 2-bottle preference test. The wild-type, *Stim1*<sup>-/-</sup>, or CdCl<sub>2</sub>-injected mice were first accustomed to water drinking in 2 bottles for a period of 24 hours. Then, the individually caged mice were allowed to choose between 0.1% colza oil emulsified in 0.3% xanthan gum in water or water with vehicle alone (0.3% xanthan gum) over a period of 12 hours. The water intake was determined by weighing the feeders.

**Release of <sup>3</sup>H-5-HT.** The release of <sup>3</sup>H-5-HT was determined according to Khan et al. (58). In brief, CD36-positive TBCs were loaded with <sup>3</sup>H-5-HT (2 mCi/3  $\times$  10<sup>8</sup> cells) for 1 hour in the incubation buffer containing (in mM): 40 Tris-HCl (pH 7.4); 25 glucose; 5 KCl; 150 NaCl; 1.5 CaCl<sub>2</sub>; 1.4 MgSO<sub>4</sub>; 1.2 ascorbate and 0.1 amino-oxyacetic acid at 37°C. At the end of incubations, the cells were washed by centrifugation (2,000 g, 5 minutes) in washing buffer (in mM): 150 NaCl, 5 KCl, 1.2 MgSO<sub>4</sub>, 1.0 ascorbate, 10 glucose, and 1.3 EGTA. Then, the cells were resuspended in release buffer containing clorgyline (100  $\mu$ M) and pargyline (100  $\mu$ M) in order to prevent 5-HT metabolism. The composition of the release buffer was identical to the washing buffer, except 2 mM CaCl<sub>2</sub> was substituted for MgSO<sub>4</sub>, EGTA was omitted, NaCl concentration was reduced to 75 mM, and cells were distributed into new tubes and then treated with LA in the presence or absence of SSO for 30 minutes. After centrifugation, the radioactivity in the supernatant and the pellet was determined. <sup>3</sup>H-5-HT release was calculated as percentage of unstimulated cells, by dividing the radioactivity of the supernatant by the radioactivity in pellet plus supernatant in each assay. The non-radioactive 5-HT was also detected by HPLC (8), and the values were corrected for the release of radioactive 5-HT.



**Statistics.** Statistical analysis of data was carried out using Statistica (version 4.1, Statsoft). Data are presented as mean  $\pm$  SEM. The significance of the differences between mean values was determined by 1-way ANOVA, followed by a least-significant-difference (LSD) test. For all the tests, the significance level chosen was  $P < 0.05$ .

**Study approval.** All study protocols involving mice were conducted as per the Declaration of Helsinki and European ethical guidelines, and the protocols were approved by the Regional Ethical Committee (Dijon, France).

## Acknowledgments

We thank J.F. Glaz, Cardiovascular Research Institute, Maastricht, The Netherlands, for the gift of SSO. This work was supported by contingent grants from the French Ministry of Higher Education and Research, and from Région Bourgogne (to N.A. Khan). We also acknowledge contingent grants from Agence Nationale pour la Recherche (ANR-07-PNRA-015 to P. Besnard). This work

was also supported by the Burgundy Council and Centre National Interprofessionnel de l'Economie Laitière (CNIEL) (HumanFATaste program to P. Besnard).

Received for publication July 13, 2011, and accepted in revised form March 7, 2012.

Address correspondence to: Naim Akhtar Khan, Physiologie de la Nutrition et Toxicologie, UMR U866 INSERM/Université de Bourgogne/AgroSup, 6 Boulevard Gabriel, F-21000 Dijon, France. Phone: 33.3.80.39.63.12; Fax: 33.3.80.39.63.30; E-mail: Naim.Khan@u-bourgogne.fr.

Naim Akhtar Khan's present address is: Physiologie de la Nutrition et Toxicologie, UMR U866 INSERM/Université de Bourgogne/AgroSup, Dijon, France.

- Chandrashekar J, Hoon MA, Ryba NJ, Zuker CS. The receptors and cells for mammalian taste. *Nature*. 2006;444(7117):288–294.
- Laugerette F, et al. CD36 involvement in orosensory detection of dietary lipids, spontaneous fat preference, and digestive secretions. *J Clin Invest*. 2005;115(11):3177–3184.
- Gaillard D, et al. The gustatory pathway is involved in CD36-mediated orosensory perception of long-chain fatty acids in the mouse. *FASEB J*. 2008;22(5):1458–1468.
- Khan NA, Besnard P. Oro-sensory perception of dietary lipids: new insights into the fat taste transduction. *Biochim Biophys Acta*. 2009;1791(3):149–155.
- Kawai T, Fushiki T. Importance of lipolysis in oral cavity for orosensory detection of fat. *Am J Physiol Regul Integr Comp Physiol*. 2003;285(2):R447–R454.
- Fukuwatari T, et al. Expression of the putative membrane fatty acid transporter (FAT) in taste buds of the circumvallate papillae in rats. *FEBS Lett*. 1997;414(2):461–464.
- Sclafani A, Ackroff K, Abumrad NA. CD36 gene deletion reduces fat preference and intake but not post-oral fat conditioning in mice. *Am J Physiol Regul Integr Comp Physiol*. 2007;293(5):R1823–R1832.
- El-Yassimi A, Hichami A, Besnard P, Khan NA. Linoleic acid induces calcium signaling, Src kinase phosphorylation, and neurotransmitter release in mouse CD36-positive gustatory cells. *J Biol Chem*. 2008;283(19):12949–12959.
- Dramane G, Akpona S, Simonin AM, Besnard P, Khan NA. Cell signaling mechanisms of gustatory perception of lipids: can the taste cells be the target of anti-obesity agents? *Curr Med Chem*. 2011;18(22):3417–3422.
- Gilbertson TA, Fontenot DT, Liu L, Zhang H, Monroe WT. Fatty acid modulation of K<sup>+</sup> channels in taste receptor cells: gustatory cues for dietary fat. *Am J Physiol*. 1997;272(4 pt 1):C1203–C1210.
- Schiffman SS, Suggs MS, Losee ML, Gatlin LA, Stagner WC, Bell RM. Effect of lipid-derived second messengers on electrophysiological taste responses in the gerbil. *Pharmacol Biochem Behav*. 1995;52(1):49–58.
- Dennis EA. Diversity of group types, regulation, and function of phospholipase A<sub>2</sub>. *J Biol Chem*. 1994;269(18):13057–13060.
- Kudo I, Murakami M. Phospholipase A<sub>2</sub> enzymes. *Prostaglandins Other Lipid Mediat*. 2002;68–69:3–58.
- Oike H, Matsumoto I, Abe K. Group IIA phospholipase A<sub>2</sub> is coexpressed with SNAP-25 in mature taste receptor cells of rat circumvallate papillae. *J Comp Neurol*. 2006;494(6):876–886.
- Harmon CM, Abumrad NA. Binding of sulfocucinimidyl fatty acids to adipocyte membrane proteins: isolation and amino-terminal sequence of an 88-kD protein implicated in transport of long-chain fatty acids. *J Membr Biol*. 1993;133(1):43–49.
- Fierro L, Lund PE, Parekh AB. Comparison of the activation of the Ca<sup>2+</sup> release-activated current I<sub>CRAC</sub> to InsP<sub>3</sub> in Jurkat T-lymphocytes, pulmonary artery endothelia and RBL-1 cells. *Pflügers Arch*. 2000;440(4):580–587.
- Lewis RS. The molecular choreography of a store-operated calcium channel. *Nature*. 2007;446(7133):284–287.
- Smani T, Zakharov SI, Csutora P, Leno E, Trepakova ES, Bolotina VM. A novel mechanism for the store-operated calcium influx pathway. *Nat Cell Biol*. 2004;6(2):113–120.
- Varga-Szabo D, et al. The calcium sensor STIM1 is an essential mediator of arterial thrombosis and ischemic brain infarction. *J Exp Med*. 2008;205(7):1583–1591.
- Baba Y, Nishida K, Fujii Y, Hirano T, Hikida M, Kurosaki T. Essential function for the calcium sensor STIM1 in mast cell activation and anaphylactic responses. *Nat Immunol*. 2008;9(1):81–88.
- Beyersdorf N, et al. STIM1-independent T cell development and effector function in vivo. *J Immunol*. 2009;182(6):3390–3397.
- Six DA, Dennis EA. The expanding superfamily of phospholipase A<sub>2</sub> enzymes: classification and characterization. *Biochim Biophys Acta*. 2000;1488(1–2):1–19.
- Singer AG, et al. Interfacial kinetic and binding properties of the complete set of human and mouse groups I, II, V, X, and XII secreted phospholipases A<sub>2</sub>. *J Biol Chem*. 2002;277(50):48535–48549.
- Murakami M, Masuda S, Kudo I. Arachidonate release and prostaglandin production by group IVC phospholipase A<sub>2</sub> (cytosolic phospholipase A<sub>2</sub>γ). *Biochem J*. 2003;372(pt 3):695–702.
- Wolf MJ, Gross RW. The calcium-dependent association and functional coupling of calmodulin with myocardial phospholipase A<sub>2</sub>. Implications for cardiac cycle-dependent alterations in phospholipolysis. *J Biol Chem*. 1996;271(35):20989–20992.
- Jenkins CM, Wolf MJ, Mancuso DJ, Gross RW. Identification of the calmodulin-binding domain of recombinant calcium-independent phospholipase A<sub>2</sub>β. Implications for structure and function. *J Biol Chem*. 2001;276(10):7129–7135.
- Mancuso DJ, Jenkins CM, Gross RW. The genomic organization, complete mRNA sequence, cloning, and expression of a novel human intracellular membrane-associated calcium-independent phospholipase A<sub>2</sub>. *J Biol Chem*. 2000;275(14):9937–9945.
- Kuda O, et al. CD36 Protein is involved in store-operated calcium flux, phospholipase A<sub>2</sub> Activation, and Production of Prostaglandin E<sub>2</sub>. *J Biol Chem*. 2011;286(20):17785–17795.
- Liu Y, et al. Laminar flow activates peroxisome proliferator-activated receptor-gamma in vascular endothelial cells. *Circulation*. 2004;110(9):1128–1133.
- Caro AA, Cederbaum AI. Role of intracellular calcium and phospholipase A<sub>2</sub> in arachidonic acid-induced toxicity in liver cells overexpressing CYP2E1. *Arch Biochem Biophys*. 2007;457(2):252–263.
- Tsumoda Y, Owayn C. Differential involvement of phospholipase A<sub>2</sub>/arachidonic acid and phospholipase C/phosphoinositol pathways during cholecystokinin receptor activated Ca<sup>2+</sup> oscillations in pancreatic acini. *Biochem Biophys Res Commun*. 1993;194(3):1194–1202.
- Törnquist K, Ekokoski E, Forss L. Thapsigargin-induced calcium entry in FRTL-5 cells: possible dependence on phospholipase A<sub>2</sub> activation. *J Cell Physiol*. 1994;160(1):40–46.
- Suh HN, Huang HT, Song CH, Lee JH, Han HJ. Linoleic acid stimulates gluconeogenesis via Ca<sup>2+</sup>/PLC, cPLA<sub>2</sub>, and PPAR pathways through GPR40 in primary cultured chicken hepatocytes. *Am J Physiol Cell Physiol*. 2008;295(6):C1518–C1527.
- Hichami A, Joshi B, Simonin AM, Khan NA. Role of three isoforms of phospholipase A<sub>2</sub> in capacitative calcium influx in human T-cells. *Eur J Biochem*. 2002;269(22):5557–5563.
- Roudbaraki MM, Drouhault R, Bacquart T, Vacher P. Arachidonic acid-induced hormone released in somatotropes: involvement of calcium. *Neuroendocrinology*. 1996;63(3):244–256.
- Mignen O, Thompson JL, Shuttleworth TJ. Ca<sup>2+</sup> selectivity and fatty acid specificity of the non-capacitative, arachidonate-regulated Ca<sup>2+</sup> (ARC) channels. *J Biol Chem*. 2003;278(12):10174–10181.
- Boittin FX, Shapovalov G, Hirn C, Rugg UT. Phospholipase A<sub>2</sub>-derived lysophosphatidylcholine triggers Ca<sup>2+</sup> entry in dystrophic skeletal muscle fibers. *Biochem Biophys Res Commun*. 2010;391(1):401–406.
- Ramdriamampita C, Tsien RY. Emptying of intracellular Ca<sup>2+</sup> stores releases a novel small messenger that stimulates Ca<sup>2+</sup> influx. *Nature*. 1993;364(6440):809–814.
- Bolotina VM, Csutora P. CIF and other mysteries of the store-operated Ca<sup>2+</sup>-entry pathway. *Trends Biochem Sci*. 2005;30(7):378–387.
- Mercer JC, et al. Large store-operated calcium selective currents due to co-expression of Orai1 or Orai2 with the intracellular calcium sensor, Stim1. *J Biol Chem*. 2006;281(34):24979–24990.
- Potier M, Trebak M. New developments in the signaling mechanisms of the store-operated calcium entry pathway. *Pflügers Arch*. 2008;457(2):405–415.
- Bolotina VM. Orai, STIM1 and iPLA<sub>2</sub> β: a view from a different perspective. *J Physiol*. 2008;586(13):3035–3042.
- Shuttleworth TJ. Arachidonic acid, ARC channels,





- and Orai proteins. *Cell Calcium*. 2009;45(6):602–610.
44. Mignen O, Thompson JL, Shuttleworth TJ. The molecular architecture of the arachidonate-regulated  $\text{Ca}^{2+}$ -selective ARC channel is a pentameric assembly of Orai1 and Orai3 subunits. *J Physiol*. 2009;587(pt 17):4181–4197.
45. Liou J, et al. STIM is a  $\text{Ca}^{2+}$  sensor essential for  $\text{Ca}^{2+}$ -store-depletion-triggered  $\text{Ca}^{2+}$  influx. *Curr Biol*. 2005;15(13):1235–1241.
46. Li Z, Lu J, Xu P, Xie X, Chen L, Xu T. Mapping the interacting domains of STIM1 and Orai1 in  $\text{Ca}^{2+}$  release-activated  $\text{Ca}^{2+}$  channel activation. *J Biol Chem*. 2007;282(40):29448–29456.
47. Csutora P, et al. Novel role for STIM1 as a trigger for calcium influx factor production. *J Biol Chem*. 2008;283(21):14524–14531.
48. Soboloff J, Spassova MA, Tang XD, Hewavitharana T, Xu W, Gill DL. Orai1 and STIM reconstitute store-operated calcium channel function. *J Biol Chem*. 2006;281(30):20661–20665.
49. Chang WC, Parekh AB. Close functional coupling between  $\text{Ca}^{2+}$  release-activated  $\text{Ca}^{2+}$  channels, arachidonic acid release, and leukotriene C4 secretion. *J Biol Chem*. 2004;279(29):29994–29999.
50. Stewart JE, Newman LP, Keast RS. Oral sensitivity to oleic acid is associated with fat intake and body mass index. *Clin Nutr*. 2011;30(6):838–844.
51. Pepino MY, Love-Gregory L, Klein S, Abumrad NA. The fatty acid translocase gene, CD36, and lingual lipase influence oral sensitivity to fat in obese subjects. *J Lipid Res*. 2012;53(3):561–566.
52. Gryniewicz G, Poenie M, Tsien RY. A new generation of  $\text{Ca}^{2+}$  indicators with greatly improved fluorescence properties. *J Biol Chem*. 1985; 260(6):3440–3450.
53. Hamill OP, Marty A, Neher E, Sakmann B, Sigworth FJ. Improved patch-clamp techniques for high resolution current recording from cells and cell-free membranes patches. *Pflugers Arch*. 1981;391(2):85–100.
54. Chouabe C, et al. Reduction of I(Ca,L) and I(to1) density in hypertrophied right ventricular cells by simulated high altitude in adult rats. *J Mol Cell Cardiol*. 1997;29(1):193–206.
55. Trepakova ES, Csutora P, Hunton DL, Marchase RB, Cohen RA, Bolotina VM. Calcium influx factor directly activates store-operated cation channels in vascular smooth muscle cells. *J Biol Chem*. 2000;275(34):26158–26163.
56. Braun A, et al. Orai1 (CRACM1) is the platelet SOC channel and essential for pathological thrombus formation. *Blood*. 2009;113(9):2056–2063.
57. Pathak N, Khandelwal S. Immunomodulatory role of piperine in cadmium induced thymic atrophy and splenomegaly in mice. *Environ Toxicol Pharmacol*. 2009;28(1):52–60.
58. Khan NA, Moulinoux JP, Deschaux P. Putrescine modulation of depolarization-induced [3H]serotonin release from fish brain synaptosomes. *Neurosci Lett*. 1996;212(1):45–48.